

KEEP OUT OF REACH OF CHILDREN

THRIVE® 12-26-26 should be stored in a cool, well-ventilated location out of direct sunlight. Spills from broken containers can be cleaned up with absorbing clay or other suitable materials. Avoid inhalation of spray mist if possible. Avoid contact with skin and eyes. Wear proper protective equipment when handling. Dispose of used containers in accordance with local, state, and federal regulations.

GENERAL INFORMATION

THRIVE® 12-26-26 provides agronomic and economic benefits in an easy to use water soluble form. Proper timing, rate, and placement is important for desired results and highly dependent on stage of crop growth, soil fertility levels, and environmental conditions. When used as a foliar fertilizer application (or leaf feeding), water soluble fertilizer concentrates (i.e. THRIVE® 12-26-26) will help maximize yields in many field, row crops, orchard, and small grain crops, and turf. Foliar application is part of an overall fertilizer strategy and should be used in conjunction with soil-applied fertilizers. THRIVE® 12-26-26 can be applied either alone, or with pesticides. However a small "jar test" is recommended to ensure compatibility of ingredients.

DIRECTIONS FOR USE

CROPS	LBS/ACRE	REMARKS*
Alfalfa	3 to 8	On alfalfa grown for seed, 3-4 applications are recommended. On alfalfa grown for hay, apply after each cutting when leaves are formed.
Apples, peaches, pears, and other fruit and nut crops.	Apply in delayed dormant, pre-bloom, and post-bloom sprays at 3 to 5 pounds per 100 gallons in dilute sprays or in concentrate sprays can be applied at 5 to 15 pounds per acre in no less than 40 to 50 gallons water per acre. Avoid combining with materials that may cause russetting.	
Blackberries, boysenberries, raspberries and other small fruit crops	As a foliar spray, use 5 to 10 pounds per acre in the spring on new growth, repeat at pre-bloom and again 3 weeks later.	
Brussels sprouts, cabbage, cauliflower and other brassica crops.	5	Apply at 7-10 day intervals using minimum 4 applications. A wetting agent may be added to ensure good coverage. Do not apply more than 1 lb. to Chinese cabbage.
Celery, lettuce, endive and other salad crops	5	Apply at 7-10 day intervals throughout the main growing season as soon as the 3 rd and 4 th leaves unfold.
Wheat, barley and rye	5-10	Begin after crop comes out of dormancy and continue at 2-3 week intervals.
Citrus	5-10	Apply at 7-10 day intervals. Use the higher rate for full bearing trees.



Thrive Completion Max



**GUARANTEED ANALYSIS
12-26-26**

Nitrogen (N) 12.0%
 4.7% Ammoniacal Nitrogen
 6.7% Nitrate Nitrogen
 0.6% Urea Nitrogen
 Available Phosphoric Acid (P₂O₅) 26.0%
 Soluble Potash (K₂O) 26.0%
 Boron (B) 0.02%
 Copper (Cu) 0.05%
 0.05% Chelated Copper
 Iron (Fe) 0.10%
 0.10% Chelated Iron
 Manganese (Mn) 0.05%
 0.05% Chelated Manganese
 Molybdenum (Mo) 0.0005%
 Zinc (Zn) 0.05%
 0.05% Chelated Zinc

Plant nutrients derived from Ammonium Phosphate, Potassium Nitrate, Urea, Potassium Phosphate, Sulfate of Potash, Sodium Borate, Sodium Molybdate, and EDTA chelates of Copper, Iron, Manganese, and Zinc. The maximum Chlorine (Cl) content is 0.25%.

Manufactured By:
Moore Ag
 11521 Excelsior Avenue
 Hanford, CA 93230
Net Weight: 25 lbs. (11.34 kg)

CROPS	LBS/ACRE	REMARKS*
Beans, peas and other vegetables	5	Begin applications after second leaf unfolds up to bloom, at 1-2 week intervals; 1-2 applications after pod formation.
Citrus (oranges, grapefruits, lemons):	Use as a pre-bloom and post-bloom spray, can be applied 3 to 5 pounds per 100 gallons of water.	
Corn	5-10	Begin when corn is approximately 8 inches high. Continue at 1-2 week intervals.
Cotton	5	Begin after formation of the second leaf for a total of 4-8 applications depending on growing conditions.
Cucumbers, melons, squash and other vine crops	5	Begin as soon as the second true leaf unfolds for a total of 6 applications.
Grapes	5-10	Apply with routine crop protection spray.
Hops	5-10	Apply 2 treatments 14 days apart when vines have grown halfway to the wires.
Peanuts	5-10	Apply in routine crop protection sprays.
Rice	5-10	Begin 3 weeks after emergence and continue at 7-14 day intervals.
Soybeans	5-10	Begin applications after second leaf unfolds up to bloom, at 1-2 week intervals; 1-2 applications after pod formation.
Sugar beets	5-10	Apply just after the fourth leaf unfolds for a total of 3 applications 2 weeks apart.
Strawberries	As a foliar spray, use 5 to 10 pounds per acre in the spring on new growth, repeat at pre-bloom and again 3 weeks later. For fall strawberries use the same spray program.	
Potatoes, tomatoes, eggplant and other solanaceous plants	5	In routine crop protection sprays at 7-10 day intervals.
Established Turf	Apply 2 to 4 oz per 1000 square feet with a water rate of 2 to 3 gallons or 10 to 20 oz per 5,000 square feet with a water rate of 10 to 15 gallons every 7 to 10 days or as needed. Shaded grass requires less plant nutrients than grass grown in full sun.	
Hydro Seeding - grass	Use 2 to 4 oz per 1000 square feet with Hydro-Seeding Slurry.	
Greenhouse crops and ornamentals	5-10	Apply in no less than 100 gallons of water as needed.
Transplanting solutions		Add 5 lbs. per 100 gallons of transplant solution.

NOTICE: Moore Ag warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Moore Ag. In no case shall Moore Ag be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer.